arXiv:1210.3411v1 [nlin.CD] 12 Oct 2012

The Information Content of Turbulence

R.T. Cerbus[1,*] and W.I. Goldburg[1]

[1]*Department of Physics and Astronomy, University of Pittsburgh, 3941 O'Hara Street, Pittsburgh PA 15221*

(Dated: October 15, 2012)

Information theory has been wonderfully successful in enhancing communication systems, but is rarely applied directly to natural phenomena. This work is one of the few attempts to treat a physical system as an information source that can be controlled experimentally. It may be the first to do so explicitly over a wide range of parameters. The system here is the velocity fluctuations in a turbulent soap film. The compressed file size of a velocity record is an estimate of the entropy and depends on the Reynolds number. The results are compared with a parallel analysis of the logistic map.

PACS numbers: 47.10.ad, 47.10.Fg, 47.27.eb,

The equations governing turbulent flows are well known, and there is no longer much doubt about their applicability. However they yield few exact results that can be compared with experiment [1]. The main reason is that a strongly turbulent velocity field implies the nonlinear coupling of a very large number of degrees of freedom, making exact solutions elusive and their implications even more so.

Rather than comparing the experimental observations with predictions based on the Navier-Stokes equations, the measurements are cast in the language of information theory. Just as the contents of a book or a photograph can be viewed as an information source that can be coded and stored in a computer, so can a velocity field. One may define the information content of such records and ask questions about this information as a function of turbulence parameters such as the Reynolds number.

There has been a significant amount of theoretical work developing the framework of communication and information [2, 3]. Information theory has also revealed interesting new ways of thinking about physics ideas ranging from Brownian motion to the uncertainty principle [4]. However, there have been few attempts to consider experimental data as a source of information. There have been even fewer such studies of turbulent flows [5–7].

This work is a study of two-dimensional turbulent velocity fluctuations in a soap film using information theory. Moreover, unlike previous work, we consider the information content of the velocity fluctuations for a wide range of Reynolds numbers. The analysis reveals a new way of thinking about turbulence as an information source and not as a solution to the Navier-Stoke's equations. The results are strikingly different than those for a parallel analysis of the logistic map.

Two questions about stored velocity information are addressed here: (a) what is the the compressibility of this information, that is, how much can its space in the computer's memory be reduced, and (b) how often do different velocity fluctuation patterns appear? A similar issue concerns the frequency of various words in texts written in typical books or newspapers. For texts written in English (or any language) one tabulates the frequency of words and then ranks the probability of their occurrence $p(k)$ in descending order. To illustrate, "the" will have rank $k = 1$ and "say" will have $k = 34$. This ranking $p(k)$ has been carried out for very many English words.

Starting with the work of Zipf [8], much attention has been devoted to the functional form of $p(k)$. The striking observation of many such ranked events (texts in various languages, city sizes, etc.) is their power law form:

$$p(k) \propto k^{-a} \tag{1}$$

with a very close to unity for several decades in k. The exponent then increases as it must for the sum of $p(k)$ to converge to unity.

Here we report similar measurements where the "words" are groups of velocity fluctuations of various magnitudes measured sequentially in time taken from a continuous velocity record $\mathbf{u}(t)$. For example, the group $(...u_i,\ u_{i+1},\ u_{i+2}, ...) = (...\text{-}34.52, \text{-}10.73, +5.49,...)$ cm/s is here regarded as a word with three letters or symbols ($L = 3$). This permits measurement of the dependence of a on L and turbulence parameters such as the Reynolds number Re. It is the initial value of the exponent of $p(k)$ that will be referred to as a.

Before addressing the information compressibility question, one must define the information content of a message, be it words, a photograph, sentences or velocity fluctuations. All of these can be coded, stored, and transmitted. We define a message as a large collection of symbols. A measure of the information or "surprise" contained in a symbol is $\log(1/p)$, where p is the probability of that symbol. The base of the logarithm depends on the number of different symbols in use and is conventionally 2 (for binary). Using the logarithm allows the definition of information to be additive when combining (multiplying) probabilities. If a symbol is very likely, such as a sunny weather forecast in the summer, then the surprise is small and the amount of information gained by receiving the symbol is also small. The quantity $\log(1/p)$ is thus small for high probability symbols. The average information of the entire message is the information entropy, usually called H

$$H = \langle \log \frac{1}{p(i)} \rangle = \sum_i p(i) \log \frac{1}{p(i)}, \qquad (2)$$

where $p(i)$ is the probability that the symbol i is encountered in a long string of symbols. This treatment of messages and many useful theorems relating to it is the work of C.E. Shannon [2].

Shannon's work forms the basis of all compression schemes used in communication theory, including the storage programs utilized by all computers. The compressibility of a message, according to the Shannon theory, depends on the probability distribution of the symbols used. A long string of 1s or 0s can be compressed to almost zero, as can a perfectly repetitious string of 0s and 1s (...101010...). The entropy here is zero, since H = -1log(1) = 0. In the opposite limit, a string of 1s and 0s, each chosen randomly by the flip of an unbiased coin, cannot be compressed at all and the entropy is a maximum. Shannon recognized that the proper measure of the degree of compressibility of a message is its entropy. Practically speaking, however, the entropy is the lower bound on the compressed length of a message [2, 3].

Shannon's theorem is valid for any coding method, but the calculated entropy and compressibility will depend on what coding is used. The true entropy must be calculated in such a way that all interdependencies of a message are included. Any correlation will reduce the information content of a message since knowing one symbol and its correlations can spoil the surprise of the next symbol. For example, it is very likely that "ver" will be followed by "y" in an English text. Likewise, using letters as the base set of symbols for coding a book yields an overestimate of the entropy since this overlooks word and sentence structures. When we only know our letters and are first beginning to read, any book looks random! Using words and correlations between words for coding will decrease the calculated entropy and increase the compressibility.

There are many algorithms that achieve close to optimal compression and so will approach the best estimate of the entropy currently possible using such a method [3, 10]. They of necessity yield message lengths slightly larger than H. For the data in the logistic map and for the laboratory-acquired data, the widely invoked Lempel-Ziv compression method is used in the form of the .zip compression algorithm [9]. We emphasize that calculating $-\sum_i p(i) \log p(i)$, where $p(i)$ is the probability of a single velocity value, has been done countless times before but is not a good estimate of the entropy.

To clarify the methods of analysis to be discussed, it is instructive to apply the same tools to a simple nonlinear map, namely the logistic map defined as

$$x_{n+1} = r[x_n(1 - x_n)], \qquad (3)$$

where r can span the interval 0 and 4. The initially chosen value x_0 is taken to lie in the interval $0 \leq x_0 \leq 1$, as do all x_n.

Since the logistic map is well known [11], it will not be discussed here other than to say that for r larger than a certain critical value, all the x_n take on many values, jumping from one x_n to the next in a seemingly random way, though of course Eq. (3) is deterministic. One is tempted to think of r in this equation as analogous to the the Reynolds number Re with $r = 4$ corresponding to $Re \to \infty$. As will be seen however, Re and r behave quite differently as control variables.

For values of r less than 3.57, almost any choice of x_0 will lead to a fixed value of x_n or produce periodic oscillations. So for this regime the logistic map is almost infinitely compressible, that is, $H \simeq 0$ and the rank $p(k)$ has no meaning. The focus here will be on slightly larger values than this.

Equation (3) is invoked for 10^6 values of n starting with randomly chosen seed values x_0. The data are analyzed by subtracting the average value $\langle x_n \rangle \simeq 0.5$ from each entry x_n and dividing by the rms value x_{rms} for all 10^6 data points. Each number is then rounded to several significant figures (in decimal units) and a short string of these data values are treated as a word. This method is chosen for continuity with the analysis of the turbulence measurements discussed below.

One finds that the Zipf exponent a for the logistic map is a slowly increasing function of the rank k, as in English texts. All curves have a similar slope, $a = 0.2$ for the first decade in k (data not shown). This slope depends on r. For $r = 3.59$, $a = 0.5$ and for $r = 3.61$, $a = 0.25$. It is notable that a is much less than unity, contrary to its value for the ranking of words in various languages, for city population sizes, earthquake magnitudes, and many other tabulations of activities created by nature and by human activity [12].

Figure 1 is a plot of the compression ratio $C(r)$ vs. r, where $C(r)$ is defined as the ratio of the file size of the compressed record to its uncompressed size. This varies between 0 and 1 and is bounded below by H (normalized). The message under compression is the 10^6 values of (normalized) x_n obtained at closely-spaced values of r in the range $3.3 \leq r \leq 4$.

For r below 3.4, $C(r)$ is near 0, as expected. This is followed by a small increase above zero after period doubling commences at $r \simeq 3.45$. Except for a few values of r, were $C(r) \leq 0.02$, the fractional amount of compression increases linearly with r in the narrow interval 3.58 $\leq r \leq 3.7$ (solid straight line in this linear plot).

Above $r \simeq 3.7$, $C(r)$ is rather constant until r reaches its maximum value of 4. In this interval $C(r)$ fluctuates about a mean value of approximately 0.17. There are several points in this range of r for which $C(r)$ is very small. These are the well known "islands of stability" that occur for the logistic map in the normally chaotic regime, where for certain values of r the system returns



FIG. 1: Compression ratio $C(r)$ versus the logistic map parameter r. After the transition to chaos, the compression (and entropy) increases. The solid line has a slope of roughly 1.1.



FIG. 2: Compression ratio C versus Reynolds number Re' for the 2D turbulent velocity data. The straight line is a logarithmic fit to the experimental data. The solid line is given by the equation $C(Re) = -0.003\log(Re) + 0.34$. The data here were saved to three significant figures before compression.

to periodicity [11].

Apparently, the chaotic regime of the logistic map is becoming increasingly random, since the entropy increases the more random a signal becomes. However, we do not understand the linear variation with r during the transition interval. The turbulence measurements will show very different behavior.

We now consider the same analysis applied to the turbulent velocity measurements in our 2D soap film flow. A soap solution is fed to the film from a reservoir and is driven by gravity. The film is drawn between two thin metal blades separated by a distance w of the order of several cm and the turbulence is generated by a comb inserted horizontally into the flow. The mean flow speed U is of the order of 100 cm/s. More experimental details can be found in previous work using this system [13].

Very long velocity time series ($> 10^6$) of the streamwise velocity component are taken with an LDV (laser Doppler velocimeter) at the lateral center of the soap film channel and interpolated for equal time spacing. The average data rate is between 1 kHz and 10 kHz.

The velocity data are then treated in the same way as the logistic map data with the mean subtracted out and the resulting difference divided by the rms value. The velocity record is then saved to several significant figures in decimal units. It is important not to have too few significant figures nor too many. Too few will lead to a coarse-grained description that throws out important details just like a grainy photograph. In the opposite limit if there are too many, statistical fluctuations are increased due to the limited length of the entire data string.

Figure 2 is a linear-log plot of C versus a Reynolds number $Re' \equiv u'w/\nu$ where w is the channel width, u' is the rms velocity (of order 10 cm/s) and ν is the kinematic viscosity (0.01 cm^2/s). If the independent variable, is taken as $Re \equiv Uw/\nu$ rather than Re', the same behavior is seen but the scatter of the data points is increased.

The compression ratio (and hence H) is clearly of logarithmic form: $C(Re) \propto -\log(Re)$. Note that $C(Re)$ is a decreasing function of the Reynolds number, in stark contrast to the logistic map (see Fig. 1), where it increases with r after turning chaotic. This is also contrary to the results of Wijesekera [6], who estimated H via the power spectra of density fluctuations in the ocean. However, it is difficult to compare given their unique definition of the Re.

This decrease in $C(Re)$ suggests an increase in the number of correlated spatial and temporal scales as Re grows [4]. There is reason to expect the opposite behavior, since the number of degrees of freedom should increase with increasing Re [14]; all else being equal a larger alphabet of symbols means a larger entropy. The decrease in $C(Re)$ is occurring in spite of the growing number of degrees of freedom that are excited as the flow becomes more turbulent.

In the higher Reynolds number range ($Re > 10^2$) the measured power spectrum of the velocity fluctuations $E(k)$ (k is the wavenumber) has a clear power law scaling with an exponent near -3 (spectral data not shown). This value is indicative of the direct enstrophy cascade usually observed in decaying 2D turbulence [13]. The power spectra measured for $Re < 10^2$ are flat, indicating that there is no self-similar scaling. It is striking that the present method of data analysis is able to capture the behavior of turbulence at Re too low for the usual

Kolmogorov scaling ideas to be applied [1].

The analysis of the probability-rank distributions can also be applied to the measurements. The distributions are plotted in Figure 3 for widely different values of Re using a word length of $L = 3$. A somewhat longer word length could have been used. To the extent that the data can be characterized by a power law exponent, it is much less than the Zipf value $a \simeq 1$.



FIG. 3: Word probability $p(k)$ versus rank k for several different Reynolds numbers, where $Re = Uw/\nu$. The word length is $L = 3$. The broadness of the distribution decreases with increasing Reynolds number. (□) Re = 49516, (○) Re = 13172, (△) Re = 1823. The dashed line is only a guide for the eye.

Since the entropy is calculated from a probability distribution, the functional form of $C(Re)$ is presumably connected with $p(k)$. Of course, these $p(k)$ are not the distributions for the coding that the compression algorithm actually uses. Note that the broadness of the distributions in Figure 3 decreases with increasing Re. This is consistent with the compression data; increasing Re increases constraints (correlations) which decreases $C(Re)$ and should narrow $p(k)$. Stated in another way, the broadness of a distribution, such as $p(k)$, is linked to the magnitude of the entropy since a completely flat distribution gives the maximum entropy.

In summary, the velocity time series of a 2D turbulent soap film is analyzed using tools from Shannon's information theory. The main result is that the compression ratio, which is an approximation to the true entropy, is measured to be a logarithmically decreasing function of the Reynolds number. As far as we know, there are no predictions for the dependence of Shannon's entropy on the Reynolds number, although the decrease is in accord with our expectations. This strikingly weak dependence of the compression ratio on the strength of the turbulence calls for a simple explanation, but so far it has eluded us.

These measurements include low Reynolds number turbulence where the cascade picture can not be applied. If the experiments were expanded to include laminar flows, the compression ratio would start out as almost zero. Assuming that the entropy is a continuous function, it must then increase with Reynolds number, just as the logistic map analyzed here and as in [6, 7]. It should reach a maximum before decreasing logarithmically in the manner of Figure 2. This slow decrease suggests that increasing the Reynolds number further may only affect the turbulence by a small amount. This is in accord with our present understanding of large Reynolds number turbulence.

Many of the tools developed under the umbrella of information theory have not yet been applied to experimental data. Shannon entropy and other measures of complexity [15] are, for the most part, waiting to be used, although there has been some work already done [5–7]. We believe the present experiments are amongst the first few steps in that direction.

We thank Mahesh Bandi for introducing us to this topic and encouraging us to think about our velocity data in this way. This work is supported by NSF Grant No. 1044105 and by the Okinawa Institute of Science and Technology (OIST). R.T.C. is supported in part by a Mellon Fellowship through the University of Pittsburgh.

---

[*] Electronic address: rtc17@pitt.edu

[1] P.A. Davidson, *Turbulence: An Introduction for Scientists and Engineers* (Oxford University Press, Oxford, 2004)
[2] C.E. Shannon, W. Weaver, *The Mathematical Theory of Communication* (University of Illinois Press, Urbana, Ill., 1964)
[3] T.M. Cover, J.A. Thomas, *Elements of Information Theory* (Wiley, New York, 1938)
[4] L. Brillouin, *Science and Information Theory* (Academic Press, New York, 1962)
[5] X.J. Wang, P. Gaspard, Phys. Rev. A **46**, R300 (1992)
[6] H.W. Wijesekera, T.M. Dillon, J. Geophys. Res. **102**, 3279 (1997)
[7] A. Brandstäter et al., Phys. Rev. Lett. **51**, 1442 (1983)
[8] G.K. Zipf, *Human Behavior and the Principle of Least Effort* (Addison-Wesley, Cambridge, Mass., 1949)
[9] D. Salomon, *Data Compression: The Complete Guide* (Springer-Verlag, London, 2007)
[10] T. Schürmann, P. Grassberger, Chaos **6**, 414 (1996)
[11] R.C. Hilborn, *Chaos and nonlinear dynamics* (Oxford University Press, New York, 1994)
[12] B. Corominas-Murtra, R.V. Solé, Physical Review E **82**, 011102 (2010)
[13] H. Kellay, W.I. Goldburg, Rep. Prog. Phys. **65**, 845-894 (2002)
[14] C.V. Tran, L. Blackburn, Physical Review E **79**, 056308 (2009)
[15] J.P. Crutchfield, Nat. Phys. **8**, 17 (2012)